

REVIEW Article

The Dual Role of Uric Acid: Pathological Implications Across Chronic Diseases and Current Approaches to Hyperuricemia Management

Aiza Zehra¹, Ayeeza Asghar¹, Sarah Ali¹, Muhammad Umair Khalid¹, Syed Imran Ali Shah²*

¹ CMH Lahore Medical University, Lahore, Pakistan

² Department of Biochemistry, Azra Naheed Medical College, Faculty of Medical Sciences, Superior University, Pakistan

* Corresponding authors: Sarah Ali, MBBS, CMH Lahore Medical College and IOD Abdur Rehman Road, Lahore Cantt, Pakistan; Email: Saraali6875@outlook.com; Phone: +923229009995.

Submitted: May 16, 2026; *Revised:* June 11, 2026; *Accepted:* June 14, 2026; *Published:* June 30, 2026.

Citation: Zehra et al. The Dual Role of Uric Acid: Pathological Implications Across Chronic Diseases and Current Approaches to Hyperuricemia Management. *Discoveries* 2026, Apr-Jun, 14(2): e231. DOI: 10.15190/d.2026.10

ABSTRACT

Uric acid is the end product of purine metabolism and plays a dichotomous role in the human body. On one hand, it exerts antioxidant and neuroprotective effects; on the other hand, chronic hyperuricemia has been strongly associated with diseases beyond gout, affecting the cardiovascular, renal, metabolic, autoimmune, and central nervous systems (CNS). Excess uric acid promotes oxidative stress, endothelial damage, neurodegeneration, inflammasome activation, and impairs energy metabolism. It exacerbates autoimmune diseases, such as Systemic Lupus Erythematosus (SLE) and antiphospholipid syndrome (APS), by increasing inflammatory and oxidative damage, leading to greater end-organ damage.

The British Society for Rheumatology, European League Against Rheumatism, American College of Rheumatology, and National Institute for Health and Care Excellence (NICE) have all established a “treat-to-target” approach for hyperuricemia with serum urate levels below 6 mg/dL and below 5 mg/dL in severe cases. Allopurinol and Febuxostat, xanthine oxidase inhibitors, are used as first-line pharmacological therapies for the treatment of hyperuricemia, whereas uricosurics and Interleukin-1

(IL-1) inhibitors are preferred in cases of refractory hyperuricemia. Lifestyle modifications, such as the Dietary Approaches to Stop Hypertension (DASH) diet, weight reduction, and smoking cessation, are also recommended for the long-term management of the disease. Sodium-Glucose Cotransporter 2 (SGLT2) inhibitors, selective urate transport inhibitors, and plant-derived anti-inflammatory compounds have emerged as new treatments with promising responses.

This review synthesizes the current literature on the multifaceted role of uric acid and emphasizes its systemic implications in chronic diseases. It also outlines the already established management options and new innovative therapies for managing this disease. Understanding this dichotomous role is essential for adopting a precise management approach that balances the protective and pathological effects of uric acid.

Keywords

Uric Acid; Hyperuricemia; Chronic Diseases; Cardiovascular Disease; Autoimmune Disorders; Therapeutic Interventions; Oxidative Stress; Chronic Kidney Disease (CKD); Inflammation.

SUMMARY

1. Introduction
2. Pathological Impact
3. Uric Acid and Chronic Diseases
4. Therapeutic Intervention
 - 4.1 Pharmacological Management (Allopurinol, Febuxostat, Uricosurics)
 - 4.2 Lifestyle and Dietary Modifications
5. Future Perspectives and Research Directions
6. Conclusion

Abbreviations

ATP-Binding Cassette Subfamily G Member 2 (ABCG2); American College of Rheumatology (ACR); Atrial Fibrillation (AF); AMP-Activated Protein Kinase – Unc-51-Like Kinase 1 Pathway (AMPK-ULK1); Antiphospholipid Syndrome (APS); Blood-Brain Barrier (BBB); British Society for Rheumatology (BSR); British Society for Rheumatology / British Health Professionals in Rheumatology (BSR/BHPR); Calcium Ion (Ca²⁺); Catalase (CAT); Chronic Kidney Disease (CKD); Cardiovascular Disease (CVD); Dietary Approaches to Stop Hypertension (DASH); European League Against Rheumatism (EULAR); Gouty Arthritis (GA); Glomerular Filtration Rate (GFR); Glucose Transporter-4 (GLUT4); Glucose Transporter-9 (GLUT9); Gasdermin D (GSDMD); Heart Failure with Reduced Ejection Fraction (HFrEF); Hyperuricemia (HUA); Immunoglobulin A (IgA); Interstitial Lung Disease (ILD); Interleukin-1 (IL-1); Interleukin-1 Beta (IL-1 β); Interleukin-6 (IL-6); Interleukin-18 (IL-18); Low-Density Lipoprotein (LDL); Malondialdehyde (MDA); Major Adverse Cardiovascular Events (MACE); Metabolic Exercise Cardiac Kidney Index (MECKI); Monosodium Urate (MSU); Myocardial Infarction (MI); National Institute for Health and Care Excellence (NICE); NLR Family Pyrin Domain-Containing 3 Inflammasome (NLRP3); Nuclear Factor Erythroid 2-Related Factor 2 (Nrf2); Nonsteroidal Anti-Inflammatory Drugs (NSAID); N-terminal pro-Brain Natriuretic Peptide (NT-proBNP); New York Heart Association (NYHA); Rheumatoid Arthritis-Associated Interstitial Lung Disease (RAILD); Reasons for Geographic and Racial Differences in Stroke Study (REGARDS); Reactive Oxygen Species (ROS); Reactive Oxygen Species–Ferroptosis (ROS-ferroptosis); Rheumatoid Arthritis (RA); Serum Uric Acid (SUA); Reactive Oxygen Species–Neutrophil Extracellular Trap (ROS-NET); Sodium-Glucose Cotransporter 2 (SGLT2); Systemic Lupus Erythematosus (SLE); Superoxide Dismutase (SOD); Toll-Like Receptor 4 (TLR4); Toll-Like Receptor 6 (TLR6); Transient Receptor Potential Melastatin-2

(TRPM2); Uric Acid (UA); Usual Interstitial Pneumonia (UIP); Urate-Lowering Therapy (ULT); Urate Transporter 1 (URAT1); Uric Acid Right for Heart Health Study (URRAH); Xanthine Oxidoreductase (XOR).

1. Introduction

Uric acid is the final product of purine breakdown in humans and has been linked to a paradoxical role in different physiological and pathological conditions. While it functions as a potent antioxidant in plasma, excessive accumulation has been implicated in numerous long-term health problems and chronic illnesses. While it is established that elevated uric acid levels have been notorious as one of the principal causes for gout and kidney stones, this paper further discusses how uric acid may also play an important role in the development of conditions like hypertension, cardiovascular disease, chronic kidney disease, autoimmune conditions, and its effects on neurological and executive function. The underlying mechanisms by which it exerts its effects are important to highlight so that therapeutic interventions can be better targeted, more effective, and tailored to the specific pathways involved in disease progression. Uric acid can cross the blood-brain barrier and accumulate in the cerebrospinal fluid, impairing neuronal function and causing cognitive decline¹. minor elevations in levels have exerted complex effects such as structural brain changes and disruptions in cognitive abilities, such as memory, concentration, and comprehension². In the cardiovascular system, hyperuricemia promotes endothelial dysfunction, atherosclerosis, and myocardial remodeling through mechanisms including oxidative stress, inflammasome activation, and disrupted energy metabolism, discussed in detail below. Through these mechanisms, hyperuricemia is invariably associated with increased risk of myocardial infarction, atrial fibrillation, and ischemia-reperfusion injury^{3–5}. Moreover, hyperuricemia has also been observed in connection with metabolic dysfunction, correlating with insulin resistance, obesity, and dyslipidemia⁶. Hyperuricemia has also been implicated in the exacerbation of autoimmune conditions, notably systemic lupus erythematosus (SLE), by amplifying oxidative and inflammatory cascades that contribute to disease progression⁷. All these findings and others, as discussed below, point to the systemic impact of hyperuricemia, beyond its classical associations.

Understanding its complex role in various disease mechanisms is crucial for the development of effective treatment strategies.

Organizations such as the British Society for Rheumatology (BSR), the European League Against Rheumatism (EULAR), the American College of Rheumatology (ACR), and the National Institute for Health and Care Excellence (NICE) have developed evidence-based guidelines and recommendations to guide the clinical management of hyperuricemia and its associated symptoms. Treatment goals focus on reducing serum urate to below the saturation threshold for crystal formation, with most authorities advocating a target of <6 mg/dL, and even lower (<5 mg/dL) in patients with severe disease manifestations⁸. Guidelines further emphasize a "treat-to-target" approach involving urate-lowering therapy (ULT) in patients with recurrent attacks, renal dysfunction, comorbidities, or evidence of joint damage⁸. First-line pharmacologic interventions include xanthine oxidase inhibitors such as allopurinol and febuxostat, while uricosuric agents

limited success, dietary patterns like the DASH and Mediterranean diets, which address underlying metabolic dysfunction, offer a more impactful approach¹⁰. Weight reduction, regular physical activity, and smoking cessation are also recommended due to their positive effects on serum urate levels¹¹.

Recent advancements in research have broadened treatment possibilities with the introduction of novel agents that act on urate transporters (URAT1), sodium glucose co-transporter 2 (SGLT2), and interleukin signaling pathways, as well as natural anti-inflammatory compounds. These recent advancements reflect a promising role in the systemic management of hyperuricemia, offering new possibilities for more effective, targeted, and individualized treatment approaches¹².

2. Pathological impact

Uric acid, a byproduct of purine breakdown, serves as both a physiological antioxidant and a pathological




Figure 1. Multisystemic Pathological Impacts of Elevated Uric Acid

This directional flow illustration maps the transition of uric acid from a passive biomarker to an active pathophysiological driver across several organ systems

and anti-inflammatory therapies provide additional options depending on patient-specific factors⁹. Beyond pharmacological management, dietary and lifestyle modifications play a crucial role in managing hyperuricemia. While low-purine diets have shown

contributor to disease progression. Excessive accumulation of uric acid, termed hyperuricemia (serum uric acid concentration >7.0 mg/dl)¹³ implicated in a wide range of disorders. These include kidney disease, neurological disease, cardiovascular

complications, as well as autoimmune conditions, revealing a complex and often contradictory role in human health¹⁴. Dysregulation of serum uric acid levels by transporters such as urate transporter 1 (URAT1), glucose transporter 9 (GLUT9), and ATP-binding cassette subfamily G member 2 (ABCG2) results in hyperuricemia. The most common acute presentation of hyperuricemia is gout, the formation of monosodium urate crystal deposits in joints and other body tissues. Understanding the dual nature of uric acid is essential to evaluating its pathological impact and therapeutic potential¹⁵.

3. Uric acid and chronic diseases

Elevated uric acid levels are increasingly associated with the development and progression of various chronic diseases. From cardiovascular and neurological disorders to kidney disease and autoimmune conditions, uric acid appears to play a complex and often detrimental role, acting as both a biomarker and a potential contributor to disease mechanisms. However, it is important to note that many of these associations are derived from observational studies, and the distinction between correlation and causation remains an active area of investigation. To better understand the role of uric acid, **Figure 1** lists the potential systems and diseases that can be involved.

Hyperuricemia may facilitate the passage of uric acid through the blood-brain barrier from the plasma into the cerebrospinal fluid (CSF), where preclinical evidence suggests it can promote neuroinflammation and neuronal impairment, potentially contributing to cognitive decline¹⁷. This is particularly important in the context of aging, where even minor fluctuations in uric acid levels may lead to anatomical changes in the brain as well as direct alterations in neural functioning, impairing active memory, concentration, understanding ability, and higher-order cognitive skills¹⁸.

There is some incongruity between elevated uric acid levels and the severity of central nervous system damage, which shows that other factors may also be involved in the underlying pathology, beyond what is traditionally understood about uric acid's role. This is corroborated by the fact that certain subtypes of dementia, such as vascular dementia occurring as a complication of underlying stroke, are more prone to hyperuricemia damage due to the damage caused to cerebral blood vessels by elevated uric acid levels¹⁹,

whereas other types of dementia, including Parkinson disease-associated dementia or Alzheimer's disease, showed no significant variation in levels between cases and controls across multiple studies². Lower serum uric acid levels are correlated with a negative disease course in Parkinson's disease. Hypouricemia has been associated with a worse clinical trajectory, characterized by decline in motor function, cognitive ability, and overall quality of life over time²⁰.

The implications of high uric acid levels in heart disease (**Table 1**) may warrant proactive measures for its management to prevent myocyte damage and lower the risk of cardiovascular disease. Table 1 outlines the various effects of high uric acid levels on the heart, along with its underlying mechanisms and resulting outcomes²¹.

The Italian URRAH (Uric Acid Right for Heart Health) study, a multicenter observational cohort (n = 23,467), proposed a prognostic cut-off of >5.7 mg/dL for predicting fatal myocardial infarction (HR 1.38, 95% CI 1.10–1.76), with the association significant in women but not men³⁵.

The advancement of coronary artery disease can also be attributed in part to uric acid levels in patients with both comorbidities as elevated uric acid may accelerate plaque disruption, calcification and the formation of hemorrhagic thrombi³⁶.

An observational study reported that for each 1 mg/dL rise in uric acid levels, the risk of atrial fibrillation was associated with a 35% increase in women compared with 15% in men³⁷. In the ARIC cohort (n = 15,382), per-SD HRs were 1.25 (95% CI 1.08–1.43) in women versus 1.05 (95% CI 0.94–1.18) in men, with a significant interaction (p < 0.01).

Piepoli et al. (2020) analyzed the MECKI database (n = 4,577 patients with heart failure with reduced ejection fraction [HFrEF]) and found serum uric acid was associated with increased total and cardiovascular mortality (HR 1.120 and 1.128, respectively; p < 0.0001). Critically, this association was significant only in New York Heart Association (NYHA) class I–II (HR 1.17, p < 0.0001) and not in class III–IV (HR 1.03, p = NS)³⁸.

Damage to heart muscle in this context has been found to correlate with elevated levels of circulatory biochemical markers, including highly sensitive troponin I and N-terminal pro-brain natriuretic peptide (NT-proBNP), further supporting an association between uric acid and deterioration of heart muscle function³⁹.

Table 1. The effects of high uric acid levels on the heart

Effects	Results	Mechanism
Direct effects on the myocardium	• Endothelial cell damage²² • Promotes atherosclerosis • Alterations in myocardial structure • Induction of exudation and fibrosis²³ • Cell shortening²⁴	• Decreases troponin affinity for calcium within cardiac myocytes
Inflammasome activation	• Cell swelling and subsequent lysis • Activated macrophages secrete regulatory factors, inducing arrhythmias in surrounding myocytes	• Increased TLR6 levels associated with NLRP3 inflammasome activation²⁵ • Production of ROS²⁶ • Caspase 1 maturation with subsequent activation of IL-18 and IL-1β and N terminal domain of Gasdermin D (GSDMD)²⁷
Disruption of Myocardial Energy Metabolism	• Cardiac hypertrophy • Accumulation of cytoplasmic lipids²⁸	• Insulin resistance in myocytes by insulin receptor substrate 1 phosphorylation, inhibiting Akt phosphorylation and GLUT4 translocation²⁹ • AMPK-ULK1 pathway stimulates autophagy³⁰ • Reduced carnitine palmitoyl transferase 1B activity affects fatty acid oxidation³¹
Diminished antioxidative potential	• Exacerbation of oxidative stress in the myocardium	• Increased levels of malondialdehyde (MDA) and decreased superoxide dismutase (SOD) and catalase (CAT) • Downregulation of nuclear factor erythroid 2-related factor 2 (Nrf2) which regulates antioxidant stress³²
Myocardial Ischemia-Reperfusion Injury	• Hypoxia/ • Reperfusion in cardiac cells by the production of reactive oxygen species and cellular proptosis³³	• Reperfusion generates oxygen radicals, activates NLRP3 inflammasome and the ROS/TRPM2/Ca²⁺ pathway³⁴

A study conducted in Bangladesh that divided participants based on their blood pressure levels showed that xanthine oxidase levels were significantly higher in hypertensive patients, encompassing even patients on anti-hypertensive medication, than in patients with normal blood pressure levels⁴⁰. through its association with hypertension, hyperuricemia has also been proposed as a potential risk factor for stroke, as suggested by the REGARDS study conducted in 2020, where hyperuricemia was associated with ischemic stroke (HR 1.40, 95% CI 1.10–1.78) after adjustment for demographic variables and blood pressure^{41,42}. Alternatively, multiple studies have failed to find an association between uric acid levels and vascular complications or mortality in patients with ischemic stroke⁴³

A cross-sectional study conducted in Japan among adolescents revealed that hyperuricemia was associated with features suggestive of metabolic syndrome, such as increased body mass index, glycosylated hemoglobin levels, non-alcoholic steatohepatitis, and increased low-density lipoprotein (LDL) levels suggestive of hypercholesterolemia⁴⁴. Xanthine oxidoreductase (XOR), an enzyme crucial for the formation of uric acid from purines, is significantly associated with insulin resistance, which stimulates its activity, providing a plausible biological explanation for why uric acid levels may be higher in patients with diabetes mellitus⁴⁵.

High uric acid levels have been implicated in the progression of several autoimmune diseases, potentially worsening their prognosis by activating inflammatory and oxidative pathways, ultimately

targeting end organs involved in the primary disease or amplifying an already toxic internal environment and further disturbing the body's equilibrium.

For example, systemic lupus erythematosus, a well-known autoimmune disease, has been observed to be complicated by hyperuricemia, as explained by the following diagram (**Figure 2**)⁷.

Nephropathy is also aggravated in diabetic patients with hyperuricemia, with levels of uric acid >420 $\mu\text{mol/l}$ for men and >360 $\mu\text{mol/l}$ for women complicating diabetic nephropathy by further reducing glomerular filtration rate (GFR), causing proteinuria and progressive renal impairment⁴⁶.

A recent cross-sectional study compared uric acid levels and its effects on patients with another autoimmune disease known as antiphospholipid syndrome (APS). It showed that the patients afflicted with APS who had higher serum uric acid levels also had greater occurrences of arterial hypertension, myocardial infarction, thrombotic microvascular disease and disordered lipid metabolism. Despite the small sample size of this study, the selection of patients with APS and no other coexisting conditions reduces confounding factors significantly and makes

hyperuricemia group had significantly worse renal outcomes (Kaplan-Meier, $p = 0.0147$), higher tubular atrophy/interstitial fibrosis scores, higher vascular injury scores, and greater glomerular sclerosis percentage (all $p < 0.01$)⁴⁸.

A meta-analysis discussing the role of uric acid in myasthenia gravis interestingly found that patients of myasthenia had lower levels of uric acid in their blood compared to healthy individuals, and more severe symptoms were found in patients with lower uric acid levels. However, the dichotomy of uric acid in myasthenia gravis lies in its dual role, while lower levels are linked to greater disease severity, its complex metabolism and potential risks at higher concentrations make it unsuitable as a direct therapeutic target⁴⁹.

Hyperuricemia has also been shown to complicate rheumatoid arthritis (RA)-associated interstitial lung disease (ILD), with higher levels in the serum and bronchoalveolar fluid of RAILD patients, specifically in the setting of usual interstitial pneumonia (UIP)⁵⁰. A larger multicenter study ($n = 829$) confirmed elevated UA as an independent risk factor for ILD in RA (OR 1.68, 95% CI 1.23–2.28) and a prognostic




Figure 2. Mechanistic Pathways of Hyperuricemia-Induced Renal Function in SLE

This schematic illuminates the synergistic pathways through which hyperuricemia aggravates renal pathology within the inflammatory baseline of SLE, linking the convergence of distinct molecular cascades to functional renal decline.

the complications mentioned above strongly attributable to high uric acid levels⁴⁷.

Lu et al. (2020) cohort studied 208 IgAN patients divided by uric acid levels and found the

indicator for mortality (HR 1.80, 95% CI 1.03–3.17)⁵¹.

The relationship between hyperuricemia and kidney disease is complex and bidirectional, as

elevated uric acid levels can both result from impaired renal function and contribute to further kidney injury through multiple pathogenic pathways. Among the renal structures affected by hyperuricemia, the tubular compartment is particularly vulnerable. Renal tubular injury may result from both urate crystal deposition and the direct effects of soluble uric acid. Monosodium urate (MSU) crystals can form within the tubular lumen when urinary uric acid concentrations exceed their solubility threshold and once deposited, these crystals may trigger inflammatory responses and tissue injury. Studies suggest that MSU crystals can activate the NLR family pyrin domain-containing 3 (NLRP3) inflammasome, a multiprotein complex that promotes maturation of pro-inflammatory cytokines such as IL-1 β and IL-18. This contributes to inflammasome activation, interstitial inflammation and fibrosis, supporting a pathogenic role for local crystal formation in kidney injury. Soluble uric acid itself can also directly contribute to renal damage by stimulating inflammatory signaling and cytokine production, including IL-1 β . In proximal tubular epithelial cells, activation of the NLRP3 inflammasome occurs through TLR4-dependent pathways and may be amplified by oxidative stress, mitochondrial dysfunction, reactive oxygen species generation, lysosomal damage, and endoplasmic reticulum stress. Collectively, these findings indicate that both crystalline and soluble forms of uric acid promote renal tubulointerstitial injury through interconnected inflammatory and oxidative mechanisms, highlighting their potential contribution to the progression of chronic kidney disease (CKD) and the complex relationship between CKD and hyperuricemia⁵².

Endothelial dysfunction is a key pathway through which hyperuricemia contributes to renal injury. Elevated uric acid levels reduce nitric oxide bioavailability by suppressing endothelial nitric oxide synthase activity while simultaneously increasing reactive oxygen species production, leading to oxidative stress⁵³. Uric acid also promotes vascular inflammation through activation of hypoxia-inducible factor-1 α (HIF-1 α), enhancing endothelial permeability and leukocyte recruitment. Together, these effects induce vascular remodeling and impair renal autoregulation, resulting in abnormal glomerular hemodynamics that may accelerate kidney injury and CKD progression⁵⁴. Lastly, hyperuricemia can alter glomerular hemodynamics

by affecting the afferent arteriole and impairing endothelial function. This may lead to either glomerular hyperfiltration or hypoperfusion. The hyperfiltration pattern is associated with proteinuria, activation of the renin–angiotensin system, and progressive glomerular injury, whereas the hypoperfusion pattern is characterized by reduced glomerular blood flow and ischemic injury with minimal proteinuria⁵⁵. All these findings suggest that elevated uric acid can contribute to CKD progression through distinct hemodynamic mechanisms.

Although uric acid is traditionally viewed as a pathogenic metabolite implicated in gout, cardiovascular disease, chronic kidney disease and metabolic disorders, emerging evidence suggests that its biological role is more complex. Uric acid exhibits a dual nature, functioning not only as a mediator of disease in increased concentrations but also as an important physiological molecule with potential protective effects. A recognized beneficial role of uric acid at normal physiological levels is its inhibition of CD38, an enzyme that breaks down nicotinamide adenine dinucleotide (NAD⁺), a molecule essential for energy production, cellular repair and immune function. By reducing CD38 activity, uric acid helps preserve NAD⁺ levels and prevent excessive inflammation. In contrast, when uric acid levels exceed the normal levels and form monosodium urate (MSU) crystals, these crystals can increase CD38 expression in immune cells, leading to greater inflammation. Therefore, while excessive uric acid can contribute to inflammatory conditions such as gout, normal levels of UA may be beneficial by maintaining NAD⁺ availability and helping regulate immune responses⁵⁶.

Uric acid also contributes significantly to the body's antioxidant defense system. Reactive oxygen species (ROS) play an essential role in regulating cellular metabolism, however excessive ROS production can damage cellular structures such as DNA, proteins, and lipids leading to cell dysfunction, cell death, and various diseases. Uric acid is recognized as an important endogenous antioxidant, and studies have demonstrated that it protects cells from oxidative damage by scavenging harmful free radicals, including peroxyl radicals, hydroxyl radicals, and singlet oxygen. It exerts this antioxidant effect by donating electrons and chelating metal ions, thereby limiting free radical reactions. During the process of ROS neutralization, uric acid is oxidized to allantoin and other metabolites, making allantoin a

useful biomarker of oxidative stress in humans. These findings highlight the important role of uric acid in protecting against oxidative stress and suggest its potential contribution to the prevention of aging, cancer, and other oxidative stress-related diseases⁵⁶.

The neuroprotective potential of uric acid has been well recognized and is primarily attributed to its ability to counter oxidative stress. Under normal physiological conditions, uric acid crosses the blood brain barrier only minimally and is produced in low amounts within the brain. However, during neurological disorders, disruption of the blood brain barrier may allow circulating uric acid to enter the central nervous system (CNS), where it can exert protective effects by scavenging reactive oxygen species. Clinical studies have reported an association between reduced uric acid levels and an increased risk of neurological diseases, while accumulating evidence suggests that uric acid may help protect against disorders such as Multiple Sclerosis (MS), Alzheimer's disease and Parkinson's disease. In multiple sclerosis, a disease characterized by demyelination in the CNS, oxidative stress is thought to play a significant role in disease progression, and experimental studies indicate that uric acid administration may help reduce disease severity through its antioxidant actions. Similarly, its precursors such as inosine have demonstrated beneficial outcomes, and clinical trials suggest that inosine supplementation may improve symptoms and disease progression in MS patients⁵⁷. Recent research has also identified increased expression of CD38 in MS, while inhibition of CD38 has been shown to elevate NAD⁺ levels, as discussed previously⁵⁸. This reduces neuroinflammation and slows disease progression. Since uric acid directly inhibits CD38, it may provide neuroprotective benefits in MS by both reducing oxidative stress and preserving the availability of NAD⁺. A study published recently also suggests that monitoring uric acid levels over time may serve as an additional biomarker for understanding treatment response in multiple sclerosis. More effective MS treatments were associated with a greater long-term increase in serum uric acid levels, suggesting that uric acid may be a useful marker of disease control and oxidative stress⁵⁹.

Several pre-clinical study findings also support a potential neuroprotective role of uric acid in Parkinson's disease. Beyond its antioxidant activity, uric acid may help slow disease progression by

modulating the Nrf2-ARE signaling pathway which regulates cellular antioxidant defenses, influencing the Akt/GSK3 β signaling pathway involved in neuronal survival and function and promoting mitochondrial remodeling and maintenance, which may protect neurons from degeneration⁶⁰. Several studies have shown an inverse relationship between uric acid levels and Parkinson's Disease risk, with higher uric acid levels being associated with a lower risk of developing the disease. In contrast, diets that elevate uric acid levels and conditions such as gout have been associated with a reduced risk of Parkinson's disease⁶¹.

Influence of Sex, Age, Ethnicity, and Comorbidity Burden on Uric Acid and Disease Outcomes

Serum uric acid levels are consistently higher in males than females during reproductive years, primarily due to the uricosuric effect of estrogen, which enhances urate excretion by upregulating intestinal ABCG2 expression through the PI3K/Akt signaling pathway⁶². However, this sex gap narrows substantially after menopause. NHANES 2007–2018 data (n = 34,144) demonstrated that while overall hyperuricemia prevalence was higher in males (21.1%) than females (17.1%), females surpassed males in both prevalence and absolute numbers after age 50–59⁶³. A longitudinal study of 8,169 Japanese women confirmed that serum urate rose sharply during perimenopause (mean increase 0.41 mg/dL by post menopause), with approximately 18% of overweight or obese women at menopause developing hyperuricemia⁶⁴.

Beyond prevalence, women with hyperuricemia appear to face disproportionately higher cardiovascular risk at equivalent uric acid levels. A German population-based study (DEGS1, n = 6,918) found that the prevalence of cardiovascular and renal diseases increased with higher uric acid levels to a much greater extent in women than in men⁶⁵. Sex-specific risk factor profiles also differ in the NHANES analysis, alcohol consumption was positively associated with hyperuricemia in males but not females, while diabetes was protective in males (OR = 0.72) but a risk factor in females (OR = 1.22). Gout prevalence increases with age, and older patients frequently present with multiple comorbidities that modify disease outcomes. Among patients with three or more cardiac-renal-metabolic conditions, hyperuricemia and poorly controlled gout

showed significant positive associations with all-cause mortality, whereas these associations were not observed in patients with gout who had achieved normal serum urate levels⁶⁶. Older patients are also more susceptible to allopurinol hypersensitivity syndrome, particularly in the setting of CKD and diuretic use, necessitating more cautious dose titration⁶⁷.

Significant ethnic disparities exist in hyperuricemia prevalence and genetic susceptibility. In the NHANES analysis, Non-Hispanic Black adults had the highest hyperuricemia prevalence (23.9% in males, 23.4% in females). These disparities are partly explained by population-specific differences in urate transporter gene variants. The ABCG2 Q141K loss-of-function variant (rs2231142) has been associated with a 1.75-fold increase in gout risk across multiple populations, with the minor allele frequency being higher in East Asian populations⁶⁸.

Ethnicity also influences treatment response. In the STOP Gout trial, non-White race was associated with significantly lower odds of achieving serum urate goal with treat-to-target therapy (aOR 0.32–0.47), independent of adherence and baseline serum urate⁶⁹. These findings underscore the need for ethnicity-informed management strategies and further investigation into the biological and social determinants of treatment disparities.

4. Therapeutic interventions

Both the British Society for Rheumatology and British Health Professionals in Rheumatology (BSR/BHPR), as well as the European League Against Rheumatism (EULAR), have issued comprehensive guidelines for the management of gout, offering clear benchmarks for clinical practice. Additionally, the National Institute for Health and Care Excellence (NICE) has provided evidence-based recommendations tailored to the UK healthcare setting, ensuring that therapeutic strategies are aligned with both international and national standards. These guidelines form the foundation for evaluating and optimizing current treatment approaches in gout⁷⁰.

One of the primary goals in the treatment of gout is to lower serum uric acid levels beyond the threshold where urate crystals can form and deposit, triggering the pathological process. Since the formation of urate is a reversible process, a serum uric acid level of below 6 mg/dL is the target⁷¹. Although

the 2017 British Society for Rheumatology (BSR) guidelines and the 2022 National Institute for health and Care Excellence (NICE) guidelines differ in their recommended target serum uric acid levels for therapy, both ultimately align in their overarching treatment approach aimed at long-term disease control and reduction of gout flares⁷². Other management strategies have been outlined by the 2021 American College of Rheumatology (ACR) and the 2016 European League Against Rheumatism (EULAR) with similar goals⁷³. **Table 2** briefly summarizes the recommendations offered by NICE, ACR, EULAR, and BSR.

4.1. Pharmacological Interventions

As mentioned previously, experts across the board and most guidelines recommend several drug classes for the management of hyperuricemia and gout. These include Xanthine Oxidase inhibitors, for example, Allopurinol and Febuxostat, that work to inhibit the enzyme Xanthine Oxidase and thus prevent the formation of uric acid cellularly⁷⁴. Uricosuric agents such as Probenecid, Dotinurad, and Benzbromarone each work to facilitate uric acid excretion from the body via different mechanisms. Some studies suggest that their usage is linked to an increased risk of acute kidney injury⁷⁵ but safer alternatives in terms of dosing, and drug options such as Dotinurad 4 mg, once daily, or Verinurad have also demonstrated lesser adverse effects and better efficacy (119). Anti-inflammatory therapy includes the use of non-steroidal anti-inflammatory (NSAID) medication, corticosteroids, and colchicine. Studies support the efficacy of NSAIDs and underscore the similar effectiveness of low-dose colchicine as compared to high-dose colchicine. Steroids may be administered orally, via the intra-articular route, or intramuscularly. Although there is a lack of placebo-controlled trials for the use of both oral and intramuscular steroids, multiple randomized controlled trials support their efficacy in the setting of acute gout⁷⁷. Amongst agents that inhibit interleukin 1, a principal mediator in the pathogenesis of gout⁷⁸. Studies have demonstrated the superiority of Canakinumab. It is an anti-interleukin 1 beta monoclonal antibody that reduces pain and inflammation in patients where the use of NSAIDs and colchicine is contraindicated⁷⁹. Alternative options such as Anakinra and Rilonacept were analyzed in a systematic review, which declared the

Table 2. A comparison and summary of NICE, ACR, EULAR, and BSR guidelines to manage hyperuricemia

Strategy	2017 BSR	2022 NICE	2021 ACR	2016 EULAR
Target serum urate level	<360 $\mu\text{mol/L}$ (6 mg/dL); lower target of <300 $\mu\text{mol/L}$ (5 mg/dL) for tophaceous or severe disease	<360 $\mu\text{mol/L}$ (6 mg/dL); lower target of <300 $\mu\text{mol/L}$ (5 mg/dL) conditional on tophi, chronic gouty arthritis, or persistent frequent flares despite <6 mg/dL	<360 $\mu\text{mol/L}$ (6 mg/dL)	<360 $\mu\text{mol/L}$ (6 mg/dL); lower target of <300 $\mu\text{mol/L}$ (5 mg/dL) particularly for severe disease (tophi, chronic symptoms, frequent flares)
Indications to start ULT in gout	Recommends discussing ULT with all patients from the first attack of gout, using a treat-to-target approach	Diagnosis of gout with a treat-to-target approach	Established gout with ≥ 2 flares/year, tophaceous gout, or radiographic damage due to gout. Conditionally recommended for first flare with CKD $\geq$ stage 3, SUA >9 mg/dL, or urolithiasis	Recurrent or complicated gout. Recommends discussing ULT close to the time of first diagnosis in patients <40 years, SUA >8 mg/dL, or with comorbidities (renal impairment, hypertension, ischemic heart disease, heart failure)
Asymptomatic hyperuricemia	ULT not recommended for asymptomatic hyperuricemia	ULT not recommended for asymptomatic hyperuricemia	Conditionally recommended against initiating ULT in asymptomatic hyperuricemia, including those with comorbid CKD, CVD, urolithiasis, or hypertension	ULT not recommended for asymptomatic hyperuricemia
Flare frequency threshold	May initiate after first or recurrent flare	Frequent flares	≥ 2 flares/year	Recurrent flares (≥ 2 /year)
Tophi/extent of damage	Tophaceous gout; chronic gouty arthritis	Chronic gouty arthritis	Tophi observed clinically or through imaging; radiographic damage due to gout	Urate-related joint disease; tophi
Renal involvement	Mentions CKD specifically (stages G3–G5); recommends lower starting dose of allopurinol in CKD	Mentions CKD stage 3–5 (G3–G5) specifically	CKD stage ≥ 3 (conditional indication for ULT after first flare)	Kidney dysfunction as a whole
Urolithiasis	History of renal stones considered an indication for ULT	Not explicitly mentioned, but falls under individuals who have experienced gout flares that do not fall into the categories mentioned above	History of urolithiasis (conditional indication for ULT after first flare)	Mentions kidney stones as an indication for ULT
Special populations requiring ULT	May initiate after first or recurrent flare; emphasizes early intervention	May initiate after first or recurrent flare	–	Young age (<40 years), SUA >8 mg/dL (480 $\mu\text{mol/L}$), and/or comorbidities: hypertension, coronary artery disease, heart failure
First-line agent in ULT	Allopurinol (start low, go slow: 100 mg/day if eGFR >60; 50 mg/day if eGFR <60; dose escalation in 50–100 mg increments monthly to target). Febuxostat as alternative if allopurinol not tolerated or contraindicated	Allopurinol OR Febuxostat. Prefer allopurinol in patients with cardiovascular disease (previous MI, stroke, unstable angina)	Allopurinol (preferred first-line, including in CKD $\geq$ stage 3). Febuxostat as alternative. Probenecid as first-line if prior two not tolerated. Emphasizes HLA-B*5801 testing in Southeast Asian and African American patients	Allopurinol is the first-line ULT for patients with normal kidney function
HLA-B5801 testing	Recommends testing prior to starting allopurinol in patients of Southeast Asian or African descent	Not explicitly addressed	Strongly recommends HLA-B5801 testing prior to allopurinol in Southeast Asian and African American patients	Recommends HLA-B*5801 testing prior to allopurinol in at-risk populations
Anti-inflammatory prophylaxis during ULT initiation	Recommends colchicine prophylaxis (up to 6 months) when starting ULT	Recommends anti-inflammatory prophylaxis	Strongly recommends concomitant anti-inflammatory prophylaxis for at least 3–6 months	Recommends prophylaxis with colchicine (first-line) for 6 months
Timing of ULT initiation	Can be started during an acute flare	Can be started during an acute flare	Recommends starting during an acute flare (with anti-inflammatory treatment)	Can be started during an acute flare

reducing the frequency of acute flares. Anakinra seems to be as effective as other treatments, though this may be influenced by how it works, the small number of studies, and differences in study designs⁸⁰.

4.2. Dietary and Lifestyle Modifications

Before any pharmacological intervention, unless indicated, dietary and lifestyle modifications are essential for the long-term management of gout⁸¹. While evidence supporting the efficacy of conventional low-purine diets in preventing hyperuricemia remains limited, poor patient adherence due to low palatability further undermines their clinical utility^{82,83}. A more pragmatic approach involves targeting modifiable metabolic factors with broader relevance. Notably, the well-documented association between hyperinsulinemia and hyperuricemia suggests that addressing insulin resistance could yield dual benefits⁸⁴. In this context, dietary interventions such as the DASH or Mediterranean diet, which are both sustainable and metabolically favorable, may serve as effective strategies for managing hyperuricemia in patients with concurrent metabolic dysfunction⁸⁵.

The Mediterranean diet emphasizes daily intake of whole grains, fruits, vegetables, legumes, nuts, and olive oil, which are rich in fiber and antioxidants. Moderate amounts of dairy (especially yogurt and cheese), fish, poultry, and eggs are included regularly, while red meat is limited to small monthly portions. Red wine in moderation (up to 2 glasses/day for men, 1 for women) is preferred for its antioxidant content. The results of three separate studies on the effectiveness of the Mediterranean diet on hyperuricemia and gout yielded promising results, with an overall decrease in serum uric acid levels⁸⁶. The DASH diet, originally designed to manage hypertension, emphasizes whole grains, fruits, vegetables, low-fat dairy, and plant-based proteins from legumes and nuts instead of animal sources⁸⁷. This diet has also produced favorable outcomes for those suffering from hyperuricemia⁸⁸.

Furthermore, lifestyle modifications such as weight reduction, smoking cessation, and overall adherence to healthy habits have been shown to lower the risk and incidence of hyperuricemia. Weight loss in overweight or obese individuals lowers serum uric acid levels by improving insulin sensitivity and enhancing renal excretion of uric acid. Regular physical activity supports gout management by

lowering inflammation, improving joint mobility, and promoting cardiovascular health. Moderate-intensity exercises like walking, swimming, or cycling are preferred, as they help prevent flares without overloading the joints. Smoking contributes to systemic inflammation and elevated cardiovascular risk, both of which can worsen gout symptoms and complicate management. Quitting smoking is strongly recommended, as it reduces flare frequency, enhances treatment effectiveness, and improves overall health outcomes^{91,89}.

5. Future Perspectives and Research Directions

Transparency in the role of UA in the development of many pathophysiological conditions has allowed for the development of targeted therapeutic interventions that help modulate UA levels in the body. While xanthine oxidase inhibitors (XOIs) such as allopurinol and febuxostat remain first-line agents for reducing serum urate levels, pharmacotherapy directed at URAT1 inhibitors, SGLT2 inhibitors, immunotherapeutics, and plant-derived compounds represents a rapidly expanding therapeutic landscape⁹⁰.

URAT1 inhibitors

Benzbromarone, a non-selective URAT1 inhibitor available in many countries but not FDA-approved due to rare hepatotoxicity, has shown cardiovascular benefit in a large Korean cohort (HR 0.82 for composite cardiovascular events vs. allopurinol; HR 0.60 for all-cause mortality⁹¹).

Lesinurad, a selective URAT1/OAT4 inhibitor, was FDA-approved in 2015 for combination use with a XOI. In RCTs, lesinurad 200 mg/day combined with a XOI achieved approximately 1 mg/dL greater serum uric acid reduction than XOI alone⁹². However, serum creatinine more than doubled in 1.8% of patients at the approved dose, and efficacy is impaired at eGFR <30 mL/min. It was subsequently withdrawn from the US market for business reasons. CYP2C9 metabolism necessitates caution with co-prescribed CYP2C9 inhibitors.

Dotinurad is a novel selective urate reabsorption inhibitor (SURI) approved in Japan. It demonstrated the highest URAT1 selectivity among available uricosurics (IC₅₀ 0.0372 μ M vs. 0.190 μ M for benzbromarone and 30.0 μ M for lesinurad), with

minimal effects on secretion transporters ABCG2 and OAT1/3⁹³. In a phase 3 trial in Chinese gout patients, dotinurad 4 mg/day was superior to febuxostat 40 mg/day in achieving serum urate ≤ 6.0 mg/dL at week 24 (73.6% vs. 38.1%; $p < 0.0001$)⁹⁴. Notably, the renal safety event incidence was only 0.5%, substantially lower than lesinurad (11.2%) and verinurad (12.0%). Exploratory data from the DIANA study suggest potential vascular benefits (improved arterial stiffness and oxidative stress markers), though confirmation in larger trials is needed.

Epaminurad is currently in phase 2b development. It demonstrated dose-dependent efficacy in a multicenter RCT of 169 gout patients: 88.89% (9 mg), 71.79% (6 mg), and 54.05% (3 mg) achieved serum urate < 6 mg/dL at week 4, compared with 0% for placebo (all $p < 0.0001$). The 9 mg response rate was comparable to the febuxostat 80 mg reference arm (84.21%), with no significant differences in adverse events versus placebo⁹⁵.

Darbinurad is another URAT1 inhibitor in clinical development, though published trial data remain limited. Other earlier-stage agents include **verinurad** (phase 2; potent efficacy but 12.0% renal safety event rate)^{96,97}, **URC-102** (phase I; dose-dependent uric acid reduction in healthy volunteers)⁹⁸, and **HP501** (phase I/IIa; up to 50% serum uric acid reduction with synergistic effects when combined with febuxostat)⁹⁹.

A model-based meta-analysis of 49 studies (10,591 participants) comparing drug classes found comparable urate-lowering efficacy between URAT1 inhibitors (37.5% reduction) and XOIs (35.4%), with URAT1 inhibitors showing lower adverse event rates (51.8% vs. 55.8%) and lower dropout rates (8% vs. 17%). However, eGFR declined by 2.5% after one year with URAT1 inhibitors versus a 0.7% increase with XOIs, highlighting the importance of renal monitoring¹⁰⁰.

SGLT2 Inhibitors

Moreover, SGLT2 inhibitors have recently shown proficiency in lowering UA levels, thereby attenuating the progression to renovascular disease¹⁰¹. SGLT2 inhibitors lower uric acid by approximately 0.6–1.5 mg/dL through transactivation of GLUT9-mediated urate efflux, URAT1 downregulation, and potential ABCG2 upregulation¹⁰². A meta-analysis of 11 RCTs confirmed significant serum uric acid

reduction (MD -0.56 mg/dL, 95% CI -0.66 to -0.46 , $p < 0.00001$)¹⁰⁰.

Uric acid, hypertension, and atrial fibrillation

When building on the hypothesis that the uricase gene mutation, which occurred in the hominids, is coupled with modern era high salt dietary habits, which can lead to earlier onset of hypertension, studies are being conducted to elucidate the effect of UA lowering agents on hypertension before the emergence of microvascular injury^{103,104}. A meta-analysis of RCTs demonstrated a favorable effect of urate-lowering treatment on systolic blood pressure (mean difference -2.55 mmHg, 95% CI -4.06 to -1.05), though large-scale outcome trials are needed¹⁰⁵. Adding to this, the strong correlation between atrial fibrillation (AF) and HUA has paved the way for studies on the role of urate-lowering therapy in decreasing the incidence of AF¹⁰⁶. A drug-target Mendelian randomization study found genetic evidence supporting URAT1 inhibition for reducing AF risk (OR 0.91, 95% Bayesian credible interval 0.85–0.97), though mediation analyses suggest the benefit may be partly indirect through reductions in hypertension (56% mediation) and heart failure (48%)¹⁰⁷.

Plant-derived compounds and immunotherapeutic approaches

More recently, natural plant-based compounds are being studied to exert anti-inflammatory effects on mice with gouty arthritis (GA), as flavonoids and carotenoids are known for their role in hampering inflammation¹⁰⁸. Moreover, usage of immunotherapeutics like inhibitors of IL-6 receptor signaling can diminish MSU-induced IL-1 β production, disrupting the progression of autoinflammatory conditions like gout¹⁰⁹. Ongoing studies also talk about targeting ROS, more specifically ROS-autophagy, ROS-NET, ROS-ferroptosis, with ROS being the common denominator for inflammation associated with GA, but their therapeutic significance is yet to be determined(9).

Major challenges and future directions

Several challenges may limit clinical translation of emerging therapies:

Renal safety: Uricosuric agents increase urinary uric acid excretion, raising nephrolithiasis risk (pooled incidence 3.35%; RR 1.63 vs. non-uricosuric

treatments)¹¹¹. Newer selective agents (e.g., dotinurad, 0.5% renal events) show markedly improved profiles compared with older agents (lesinurad, 11.2%)¹¹². Combination with XOIs can reduce renal uric acid excretion by 48%, mitigating this risk.

Drug-drug interactions: CYP2C9 metabolism (lesinurad) and non-selective transporter inhibition (benzbromarone) pose interaction risks in the polypharmacy settings common among gout patients. Newer selective agents appear to have minimal transporter effects.

Patient selection: Uricosuric efficacy depends on adequate renal function (most not recommended at eGFR <30 mL/min). Phenotyping patients as urate underexcretors versus overproducers could guide therapy selection but is not routine practice.

Long-term outcomes and cost-effectiveness: Long-term data on gout flare reduction, tophus resolution, cardiovascular events, and renal preservation remain limited for most newer agents. Urate-lowering therapy has not yet demonstrated consistent cardiovascular benefit in RCTs (OR 0.67, 95% CI 0.44–1.03 for major adverse cardiovascular events [MACE]). Comparative cost-effectiveness analyses will be essential as newer agents enter the market.

Future research should prioritize large-scale RCTs evaluating long-term clinical outcomes of emerging therapies, the development of dual-target XOR/URAT1 inhibitors, integration of SGLT2 inhibitors into gout management algorithms, and translation of preclinical findings on plant-derived and ROS-targeted therapies into human studies.

6. Conclusion

This review has demonstrated the dual role of uric acid as both a protective and pathogenic entity across multiple organ systems. At physiological concentrations, uric acid functions as a potent endogenous antioxidant. It scavenges peroxyl radicals, hydroxyl radicals, and singlet oxygen and contributes to immune regulation through inhibition of CD38 and preservation of NAD⁺ availability. However, when levels exceed the physiological threshold, hyperuricemia promotes inflammation, endothelial dysfunction, oxidative stress, and end-

organ damage through pathways involving NLRP3 inflammasome activation, nitric oxide depletion, and crystal-mediated tissue injury.

In the cardiovascular system, hyperuricemia has been associated with endothelial damage, atherosclerosis, myocardial remodeling, and arrhythmogenesis through mechanisms including inflammasome activation, disruption of myocardial energy metabolism, and diminished antioxidative potential. In the kidney, both crystalline and soluble forms of uric acid promote tubulointerstitial injury through interconnected inflammatory and oxidative mechanisms, with endothelial dysfunction and altered glomerular hemodynamics contributing to CKD progression. Hyperuricemia has also been implicated in the exacerbation of autoimmune conditions, including systemic lupus erythematosus, antiphospholipid syndrome, IgA nephropathy, and rheumatoid arthritis-associated interstitial lung disease, by amplifying inflammatory and oxidative cascades that target end organs. In the metabolic domain, the association between xanthine oxidoreductase activity and insulin resistance provides a plausible biological link between hyperuricemia and metabolic syndrome.

The neurological implications of uric acid are particularly noteworthy for their paradoxical nature. Hyperuricemia facilitates the passage of uric acid through the blood-brain barrier and promote neuroinflammation and cognitive decline, particularly in the context of vascular dementia, but at the same time, lower serum uric acid levels have been consistently associated with worse outcomes in neurodegenerative diseases. In Parkinson's disease, preclinical evidence suggests that uric acid may slow disease progression by modulating the Nrf2-ARE signaling pathway, influencing the Akt/GSK3 β pathway involved in neuronal survival, and promoting mitochondrial remodeling. Several epidemiological studies have demonstrated an inverse relationship between uric acid levels and Parkinson's disease risk, with higher levels associated with reduced disease incidence. Similarly, in multiple sclerosis, uric acid may confer neuroprotective benefits through both its antioxidant actions and its inhibition of CD38, thereby preserving NAD⁺ availability and reducing neuroinflammation. Recent longitudinal data further suggest that serum uric acid trajectories may serve as a biomarker for treatment response in MS. This duality, where uric acid is protective in the central nervous system yet harmful

Major Discoveries

- Uric acid functions as both an antioxidant and a pro-inflammatory agent depending on its concentration.
- Hyperuricemia is implicated in cardiovascular disease, metabolic syndrome, renal dysfunction, and autoimmune disorders beyond gout.
- Emerging therapies target uric acid transporters and SGLT2 inhibitors, showing promise not only in reducing serum urate but also in preventing complications like stroke and renovascular disease.
- UA-lowering therapy can be considered for early hypertension and atrial fibrillation prevention.
- New plant-based anti-inflammatory compounds, IL-6 receptor blockers, and reactive oxygen species (ROS)-targeting therapies are being investigated to disrupt gout-related inflammation.

Open Questions

- Is there a threshold serum uric acid (SUA) level below which antioxidant benefits outweigh pathological risks?
- What thresholds of uric acid elevation consistently predict disease onset across populations?
- Can precision medicine approaches help tailor UA-lowering therapy based on genetic or metabolic profiles?
- What is the long-term efficacy and safety of novel pharmacological agents (e.g., DOTINURAD, IL-1 inhibitors) in large, diverse populations?
- Could plant-derived anti-inflammatory compounds be integrated into mainstream therapy for hyperuricemia?

in the cardiovascular and renal systems, represents one of the most clinically significant and unresolved aspects of uric acid biology.

The clinical management of hyperuricemia is guided by international consensus, with the BSR, EULAR, ACR, and NICE guidelines. All of these guidelines advocate a treat-to-target approach with serum urate levels below 6 mg/dL and below 5 mg/dL in severe disease. Xanthine oxidase inhibitors remain first-line urate-lowering therapy. Lifestyle modifications including the DASH and Mediterranean diets, weight reduction, and smoking cessation also serve as important adjunctive strategies. The therapeutic options are rapidly expanding with newer selective URAT1 inhibitors such as dotinurad and epaminurad demonstrating promising efficacy and improved renal safety profiles, and SGLT2 inhibitors offering urate-lowering effects alongside established cardioprotective and nephroprotective benefits. Preclinical studies on plant-derived anti-inflammatory compounds, including flavonoids and carotenoids, have shown potential in modulating gout-related inflammation through inhibition of NF- κ B activation and NLRP3 inflammasome components. Separately, biologic agents targeting key inflammatory cytokines, such as IL-1 inhibitors (anakinra, canakinumab) and IL-6 receptor antagonists (tocilizumab), represent distinct immunotherapeutic strategies for crystal-induced

arthritis refractory to conventional treatment. Additionally, ROS-targeted therapies addressing interconnected pathways of autophagy, NETosis, and ferroptosis are gaining momentum, though their clinical significance remains to be established.

Despite these advances, several important limitations and unresolved questions warrant consideration. First, many of the associations between hyperuricemia and chronic diseases discussed in this review are derived from observational studies, and the distinction between correlation and causation remains an active area of investigation. Establishing definitive causality will require well-designed prospective studies and Mendelian randomization analyses. Second, the optimal serum uric acid threshold that balances antioxidant benefits against pathological risks has not been established, and it remains unclear whether this threshold varies across disease states and patient populations. Third, long-term data on gout flare reduction, tophus resolution, cardiovascular events, and renal preservation remain limited for most newer pharmacological agents, and urate-lowering therapy has not yet demonstrated consistent cardiovascular benefit in randomized controlled trials. Fourth, the conflicting roles of uric acid across different organ systems raise fundamental questions about whether systemic urate-lowering strategies may inadvertently compromise neuroprotective effects while addressing cardiovascular and renal risk.

Future research should prioritize large-scale randomized controlled trials evaluating long-term clinical outcomes of emerging therapies, the development of dual-target XOR/URAT1 inhibitors, integration of SGLT2 inhibitors into gout management algorithms, and translation of preclinical findings on plant-derived and ROS-targeted therapies into human studies. Precision medicine approaches that tailor urate-lowering therapy based on genetic, metabolic, and disease-specific profiles may ultimately be necessary to navigate the complex and often contradictory biological effects of uric acid across the spectrum of chronic disease.

Acknowledgements

This research received no grants from any funding agency in the public, commercial or non-profit sector. Conceptualization: Aiza Zehra, Ayeeza Asghar, Sarah Ali, Muhammad Umair Khalid. Data curation: Aiza Zehra, Ayeeza Asghar, Sarah Ali, Muhammad Umair Khalid. Project administration: Aiza Zehra, Ayeeza Asghar. Supervision: Syed Imran Ali Shah. Visualization: Aiza Zehra, Ayeeza Asghar. Writing – original draft: Aiza Zehra, Ayeeza Asghar. Writing – review & editing: Aiza Zehra, Muhammad Umair Khalid, Sarah Ali.

Conflict of Interest

The author has no conflicts of interest to disclose.

Declaration of Generative AI and AI-assisted technologies

In the course of preparing this work, the author(s) utilized ChatGPT and Gemini to enhance language clarity, grammar, and conciseness. These tools were employed solely for language enhancement, and not for generating any of the scientific ideas, interpretations, analysis, or conclusions presented in this paper. After using these tools, the author(s) reviewed and edited the paper as needed, and therefore, take full responsibility for the final content.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Alam AB, Wu A, Power MC, West NA, Alonso A. Associations of serum uric acid with incident dementia and cognitive decline in the ARIC-NCS cohort. *J Neurol Sci.* 2020;414. doi:10.1016/j.jns.2020.116866
2. Khan AA, Quinn TJ, Hewitt J, Fan Y, Dawson J. Serum uric acid level and association with cognitive impairment and dementia: systematic review and meta-analysis. *Age (Omaha).* 2016;38(1):16. doi:10.1007/S11357-016-9871-8
3. Sun X, Jiao H, Zhao J, Wang X, Lin H. Rule of UA on Cardiac Myocytes Uric Acid Differently Influence the Oxidative Damage Induced by Acute Exposure of High Level of Glucose in Chicken Cardiac Myocytes. *Front Vet Sci.* 2020;7:602419. doi:10.3389/FVETS.2020.602419/BIBTEX
4. Wu H, Dai R, Wang M, Chen C. Uric acid promotes myocardial infarction injury via activating pyrin domain-containing 3 inflammasome and reactive oxygen species/transient receptor potential melastatin 2/Ca²⁺ pathway. *BMC Cardiovasc Disord.* 2023;23(1):1-13. doi:10.1186/S12872-023-03040-1/FIGURES/6
5. Xiong J, Shao W, Yu P, et al. Hyperuricemia Is Associated With the Risk of Atrial Fibrillation Independent of Sex: A Dose-Response Meta-Analysis. *Front Cardiovasc Med.* 2022;9:865036. doi:10.3389/FCVM.2022.865036/FULL
6. Yanai H, Adachi H, Hakoshima M, Katsuyama H. Molecular Biological and Clinical Understanding of the Pathophysiology and Treatments of Hyperuricemia and Its Association with Metabolic Syndrome, Cardiovascular Diseases and Chronic Kidney Disease. *Int J Mol Sci.* 2021;22(17):9221. doi:10.3390/IJMS22179221
7. dos Santos M, Veronese FV, Moresco RN. Uric acid and kidney damage in systemic lupus erythematosus. *Clinica Chimica Acta.* 2020;508:197-205. doi:10.1016/j.cca.2020.05.034
8. Kim WJ, Song JS, Choi ST. The Role of a “Treat-to-Target” Approach in the Long-Term Renal Outcomes of Patients with Gout. *J Clin Med.* 2019;8(7):1067. doi:10.3390/JCM8071067
9. Borghi C, Domienik-Karłowicz J, Tykarski A, et al. Expert consensus for the diagnosis and treatment of patients with hyperuricemia and high cardiovascular risk: 2023 update. *Cardiol J.* 2024;31(1):1. doi:10.5603/CJ.98254
10. Juraschek SP, Yokose C, McCormick N, Miller ER, Appel LJ, Choi HK. Effects of Dietary Patterns on Serum Urate: Results From a Randomized Trial of the Effects of Diet on Hypertension. *Arthritis &*

- Rheumatology. 2021;73(6):1014-1020. doi:10.1002/art.41614
11. Yokose C, McCormick N, Choi HK. Dietary and Lifestyle-Centered Approach in Gout Care and Prevention. *Curr Rheumatol Rep*. 2021;23(7):51. doi:10.1007/s11926-021-01020-y
12. Kochanowska A, Rusztyn P, Szczerkowska K, et al. Sodium-Glucose Cotransporter 2 Inhibitors to Decrease the Uric Acid Concentration—A Novel Mechanism of Action. *J Cardiovasc Dev Dis*. 2023;10(7):268. doi:10.3390/jcdd10070268
13. Leslie SW, Minter DA. Hyperuricemia. 2026.
14. Wen S, Arakawa H, Tamai I. Uric acid in health and disease: From physiological functions to pathogenic mechanisms. *Pharmacol Ther*. 2024;256:108615. doi:10.1016/J.PHARMTHERA.2024.108615
15. Dalbeth N, Gosling AL, Gaffo A, Abhishek A. Gout. *The Lancet*. 2021;397(10287):1843-1855. doi:10.1016/S0140-6736(21)00569-9
16. Du L, Zong Y, Li H, et al. Hyperuricemia and its related diseases: mechanisms and advances in therapy. *Signal Transduct Target Ther*. 2024;9(1):212. doi:10.1038/s41392-024-01916-y
17. Mijailovic NR, Vesic K, Borovcanin MM. The Influence of Serum Uric Acid on the Brain and Cognitive Dysfunction. *Front Psychiatry*. 2022;13:828476. doi:10.3389/FPSYT.2022.828476
18. Lin L, Zheng LJ, Joseph Schoepf U, et al. Uric acid has different effects on spontaneous brain activities of males and females: A cross-sectional resting-state functional MR imaging study. *Front Neurosci*. 2019;13(JUL). doi:10.3389/FNINS.2019.00763,
19. Ruggiero C, Cherubini A, Lauretani F, et al. Uric Acid and Dementia in Community-Dwelling Older Persons. *Dement Geriatr Cogn Disord*. 2009;27(4):382. doi:10.1159/000210040
20. Seifar F, Dinasarapu AR, Jinnah HA. Uric Acid in Parkinson Disease: What is the Connection? *Mov Disord*. 2022;37(11):2173. doi:10.1002/MDS.29209
21. Zheng Y, Chen Z, Yang J, et al. The Role of Hyperuricemia in Cardiac Diseases: Evidence, Controversies, and Therapeutic Strategies. *Biomolecules*. 2024;14(7). doi:10.3390/BIOM14070753,
22. Zhang J, Lin X, Xu J, Tang F, Tan L. CTRP3 protects against uric acid-induced endothelial injury by inhibiting inflammation and oxidase stress in rats. *Exp Biol Med*. 2021;247(2):174. doi:10.1177/15353702211047183
23. Lin C, Zheng Q, Li Y, et al. Assessment of the influence on left ventricle by potassium oxonate and hypoxanthine-induced chronic hyperuricemia. *Exp Biol Med*. 2022;248(2):165. doi:10.1177/15353702221120113
24. Weber M, Schreckenberger R, Schlüter KD. Uric Acid Deteriorates Load-Free Cell Shortening of Cultured Adult Rat Ventricular Cardiomyocytes via Stimulation of Arginine Turnover. *Biology (Basel)*. 2022;12(1):4. doi:10.3390/BIOLOGY12010004
25. Li P, Kurata Y, Taufiq F, et al. Kv1.5 channel mediates monosodium urate-induced activation of NLRP3 inflammasome in macrophages and arrhythmogenic effects of urate on cardiomyocytes. *Mol Biol Rep*. 2022;49(7):5939. doi:10.1007/S11033-022-07378-1
26. Wu H, Dai R, Wang M, Chen C. Uric acid promotes myocardial infarction injury via activating pyrin domain-containing 3 inflammasome and reactive oxygen species/transient receptor potential melastatin 2/Ca2+pathway. *BMC Cardiovasc Disord*. 2023;23(1):10. doi:10.1186/S12872-023-03040-1
27. Liao Y, Liu K, Zhu L. Emerging Roles of Inflammasomes in Cardiovascular Diseases. *Front Immunol*. 2022;13. doi:10.3389/FIMMU.2022.834289,
28. Yang Y, Lin C, Zheng Q, et al. L-carnitine attenuated hyperuricemia-associated left ventricular remodeling through ameliorating cardiomyocytic lipid deposition. *Front Pharmacol*. 2023;14:1016633. doi:10.3389/FPHAR.2023.1016633/BIBTEX
29. Jiao Z, Chen Y, Xie Y, Li Y, Li Z. Metformin protects against insulin resistance induced by high uric acid in cardiomyocytes via AMPK signalling pathways in vitro and in vivo. *J Cell Mol Med*. 2021;25(14):6733-6745. doi:10.1111/JCMM.16677;SUBPAGE:STRING:FUL L
30. Hardie DG. AMPK: a key regulator of energy balance in the single cell and the whole organism. *Int J Obes*. 2008;32(S4):S7-S12. doi:10.1038/ijo.2008.116
31. Yang Y, Lin C, Zheng Q, et al. L-carnitine attenuated hyperuricemia-associated left ventricular remodeling through ameliorating cardiomyocytic lipid deposition. *Front Pharmacol*. 2023;14. doi:10.3389/fphar.2023.1016633
32. Sun X, Jiao H, Zhao J, Wang X, Lin H. Rule of UA on Cardiac Myocytes Uric Acid Differently Influence the Oxidative Damage Induced by Acute Exposure of High Level of Glucose in Chicken Cardiac Myocytes. *Front Vet Sci*. 2020;7. doi:10.3389/fvets.2020.602419
33. Shen S, He F, Cheng C, Xu B, Sheng J. Uric acid aggravates myocardial ischemia-reperfusion injury via ROS/NLRP3 pyroptosis pathway. *Biomedicine &*

- Pharmacotherapy. 2021;133:110990. doi:10.1016/j.biopha.2020.110990
34. Wu H, Dai R, Wang M, Chen C. Uric acid promotes myocardial infarction injury via activating pyrin domain-containing 3 inflammasome and reactive oxygen species/transient receptor potential melastatin 2/Ca²⁺ pathway. *BMC Cardiovasc Disord.* 2023;23(1):10. doi:10.1186/s12872-023-03040-1
35. Maloberti A, Mengozzi A, Russo E, et al. The Results of the URRAH (Uric Acid Right for Heart Health) Project: A Focus on Hyperuricemia in Relation to Cardiovascular and Kidney Disease and its Role in Metabolic Dysregulation. *High Blood Pressure & Cardiovascular Prevention.* 2023;30(5):411-425. doi:10.1007/s40292-023-00602-4
36. Kobayashi N, Hata N, Tsurumi M, et al. Relation of Coronary Culprit Lesion Morphology Determined by Optical Coherence Tomography and Cardiac Outcomes to Serum Uric Acid Levels in Patients With Acute Coronary Syndrome. *Am J Cardiol.* 2018;122(1):17-25. doi:10.1016/j.amjcard.2018.03.022
37. Xiong J, Shao W, Yu P, et al. Hyperuricemia Is Associated With the Risk of Atrial Fibrillation Independent of Sex: A Dose-Response Meta-Analysis. *Front Cardiovasc Med.* 2022;9. doi:10.3389/fcvm.2022.865036
38. Piepoli MF, Salvioni E, Corrà U, et al. Increased serum uric acid level predicts poor prognosis in mildly severe chronic heart failure with reduced ejection fraction. An analysis from the MECKI score research group. *Eur J Intern Med.* 2020;72:47-52. doi:10.1016/j.ejim.2019.11.003
39. Carnicelli AP, Sun JL, Alhanti B, et al. Elevated Uric Acid Prevalence and Clinical Outcomes in Patients with Heart Failure with Preserved Ejection Fraction: Insights from RELAX. *Am J Med.* 2020;133(12):e716-e721. doi:10.1016/j.amjmed.2020.03.054
40. Miah R, Fariha KA, Sony SA, et al. Association of serum xanthine oxidase levels with hypertension: a study on Bangladeshi adults. *Sci Rep.* 2022;12(1):21727. doi:10.1038/s41598-022-26341-5
41. Chaudhary NS, Gaffo AL, Cushman M, et al. Abstract 188: Association Between Uric Acid and Stroke in the REGards Case-cohort Study. *Stroke.* 2019;50(Suppl_1). doi:10.1161/str.50.suppl_1.188
42. Chaudhary K, Malhotra K, Sowers J, Aroor A. Uric Acid - Key Ingredient in the Recipe for Cardiorenal Metabolic Syndrome. *Cardiorenal Med.* 2013;3(3):208-220. doi:10.1159/000355405
43. Zhang M, Wang Y, Wang K, Yin R, Pan X, Ma A. Association between uric acid and the prognosis of acute ischemic stroke: A systematic review and meta-analysis. *Nutrition, Metabolism and Cardiovascular Diseases.* 2021;31(11):3016-3023. doi:10.1016/j.numecd.2021.07.031
44. Aoki Y, Sofue T, Kawakami R, et al. Prevalence and factors related to hypouricemia and hyperuricemia in schoolchildren: results of a large-scale cross-sectional population-based study conducted in Japan. *Sci Rep.* 2022;12(1):17848. doi:10.1038/s41598-022-19724-1
45. Kurajoh M, Fukumoto S, Murase T, et al. Insulin Resistance Associated with Plasma Xanthine Oxidoreductase Activity Independent of Visceral Adiposity and Adiponectin Level: MedCity21 Health Examination Registry. *Int J Endocrinol.* 2019;2019. doi:10.1155/2019/1762161,
46. Lee WL, Wang PH. Uric acid in diabetic nephropathy. *Journal of the Chinese Medical Association.* 2020;83(8):794. doi:10.1097/JCMA.0000000000000349,
47. Carvalho JF, Shoenfeld Y. Uric acid levels in primary antiphospholipid syndrome. *Lupus.* 2021;30(13):2176-2178. doi:10.1177/09612033211045922,
48. Qin A, Yang D, Wang S, et al. Uric acid-based ratios for predicting renal failure in Chinese IgA nephropathy patients. *Int J Med Sci.* 2023;20(12):1584. doi:10.7150/IJMS.85430
49. Shen Y, Pang L, Wang H, et al. Comprehensive Analysis of Uric Acid and Myasthenia Gravis: IGF1R as a Protective Factor and Potential Therapeutic Target. *CNS Neurosci Ther.* 2025;31(3):e70361. doi:10.1111/CNS.70361
50. Wang Z, Wang W, Xiang T, Gong B, Xie J. Serum Uric Acid as a Diagnostic Biomarker for Rheumatoid Arthritis-Associated Interstitial Lung Disease. *Inflammation.* 2022;45(4):1800. doi:10.1007/S10753-022-01661-W
51. Hu Q, Cheng X, Lan Y, Lei H, Niu C, Luo Y. Serum uric acid predictive value and prognostic impact in rheumatoid arthritis' associated interstitial lung disease. *Front Med (Lausanne).* 2025;12:1623557. doi:10.3389/fmed.2025.1623557
52. Takata T, Mae Y, Hoi S, Iyama T, Isomoto H. Hyperuricemia in Chronic Kidney Disease: Emerging Pathophysiology and a Novel Therapeutic Strategy. *Int J Mol Sci.* 2025;26(18):9000. doi:10.3390/IJMS26189000
53. Dissanayake L V., Spires DR, Palygin O, Staruschenko A. Effects of uric acid dysregulation on the kidney. *Am J Physiol Renal Physiol.*

- 2020;318(5):F1252-F1257.
doi:10.1152/ajprenal.00066.2020
54. Jung SW, Kim SM, Kim YG, Lee SH, Moon JY. Uric acid and inflammation in kidney disease. *Am J Physiol Renal Physiol.* 2020;318(6):F1327-F1340. doi:10.1152/AJPRENAL.00272.2019
55. Takata T, Mae Y, Hoi S, Iyama T, Isomoto H. Hyperuricemia in Chronic Kidney Disease: Emerging Pathophysiology and a Novel Therapeutic Strategy. *Int J Mol Sci.* 2025;26(18):9000. doi:10.3390/IJMS26189000
56. Wen S, Arakawa H, Tamai I. Uric acid in health and disease: From physiological functions to pathogenic mechanisms. *Pharmacol Ther.* 2024;256. doi:10.1016/j.pharmthera.2024.108615
57. Gherghina ME, Peride I, Tiglis M, Neagu TP, Niculae A, Checherita IA. Uric Acid and Oxidative Stress—Relationship with Cardiovascular, Metabolic, and Renal Impairment. *Int J Mol Sci.* 2022;23(6):3188. doi:10.3390/IJMS23063188
58. Wen S, Arakawa H, Yokoyama S, Shirasaka Y, Higashida H, Tamai I. Functional identification of soluble uric acid as an endogenous inhibitor of CD38. Published online June 6, 2023. doi:10.1101/2023.06.03.543541
59. Cucu LE, Săcărescu A, Oancea A, et al. Serum Uric Acid Trajectories in Multiple Sclerosis: A 5-Year Longitudinal Comparison of High- and Moderate-Efficacy Therapies. *Med Sci (Basel).* 2026;14(2):256. doi:10.3390/MEDSCI14020256/S1
60. Xu L, Li C, Wan T, et al. Targeting uric acid: a promising intervention against oxidative stress and neuroinflammation in neurodegenerative diseases. *Cell Commun Signal.* 2025;23(1):4. doi:10.1186/S12964-024-01965-4
61. Crotty GF, Ascherio A, Schwarzschild MA. Targeting urate to reduce oxidative stress in Parkinson disease. *Exp Neurol.* 2017;298(Pt B):210. doi:10.1016/J.EXPNEUROL.2017.06.017
62. Liu L, Zhao T, Shan L, Cao L, Zhu X, Xue Y. Estradiol regulates intestinal ABCG2 to promote urate excretion via the PI3K/Akt pathway. *Nutr Metab (Lond).* 2021;18(1):63. doi:10.1186/s12986-021-00583-y
63. Zou Y, Zhang L, Xu J, et al. Sex, age, and racial/ethnic disparities in hyperuricemia prevalence and risk factors among U.S. adults: An analysis of NHANES 2007-2018 data. *PLoS One.* 2026;21(2):e0337714. doi:10.1371/journal.pone.0337714
64. Billa S, Fukui S, Paudel ML, et al. Longitudinal Changes in Serum Urate Levels From Premenopause Through Postmenopause: Interrupted Time-Series Analyses. *Arthritis Rheumatol.* 2026;78(3):724-733. doi:10.1002/art.43406
65. Engel B, Hoffmann F, Freitag MH, Jacobs H. Should we be more aware of gender aspects in hyperuricemia? Analysis of the population-based German health interview and examination survey for adults (DEGS1). *Maturitas.* 2021;153:33-40. doi:10.1016/j.maturitas.2021.08.002
66. Han Y, Yao M, Zhao H, et al. Exploration of the Interrelationship Between Serum Uric Acid, Gout, and Cardiac, Renal, and Metabolic Conditions in Middle Aged and Older People. *J Am Heart Assoc.* 2025;14(7):e038723. doi:10.1161/JAHA.124.038723
67. Kang EH. Considerations for Choosing First-Line Urate-Lowering Treatment in Older Patients with Comorbid Conditions. *Drugs Aging.* 2022;39(12):923-933. doi:10.1007/s40266-022-00986-3
68. Zhang L, Spencer KL, Voruganti VS, et al. Association of functional polymorphism rs2231142 (Q141K) in the ABCG2 gene with serum uric acid and gout in 4 US populations: the PAGE Study. *Am J Epidemiol.* 2013;177(9):923-932. doi:10.1093/aje/kws330
69. Helget LN, O'Dell JR, Newcomb JA, et al. Determinants of Achieving Serum Urate Goal with Treat-to-Target Urate-Lowering Therapy in Gout. *Arthritis Rheumatol.* 2024;76(4):638-646. doi:10.1002/art.42731
70. Mei Y, Dong B, Geng Z, Xu L. Excess Uric Acid Induces Gouty Nephropathy Through Crystal Formation: A Review of Recent Insights. *Front Endocrinol (Lausanne). Frontiers Media S.A.* 2022;13. doi:10.3389/fendo.2022.911968
71. Mackenzie IS, Ford I, Nuki G, et al. Long-term cardiovascular safety of febuxostat compared with allopurinol in patients with gout (FAST): a multicentre, prospective, randomised, open-label, non-inferiority trial. *The Lancet.* 2020;396:1745-1757. doi:10.1016/S0140-6736(20)32234-0
72. Gout: diagnosis and management. <https://www.nice.org.uk/guidance/ng219>.
73. Richette P, Doherty M, Pascual E, et al. 2016 updated EULAR evidence-based recommendations for the management of gout. *Ann Rheum Dis. BMJ Publishing Group.* 2017;76:29-42. doi:10.1136/annrheumdis-2016-209707
74. Kraev KI, Geneva-Popova MG, Hristov BK, et al. Celebrating Versatility: Febuxostat's Multifaceted Therapeutic Application. *Life (Basel).* 2023;13. doi:10.3390/life13112199

75. Hill-McManus D, Soto E, Marshall S, Lane S, Hughes D. Impact of non-adherence on the safety and efficacy of uric acid-lowering therapies in the treatment of gout. *Br J Clin Pharmacol*. 2018;84:142-152. doi:10.1111/bcp.13427
76. Li YJ, Chen LR, Yang ZL, et al. Comparative efficacy and safety of uricosuric agents in the treatment of gout or hyperuricemia: a systematic review and network meta-analysis. *Clin Rheumatol*. 2023;42:215-224. doi:10.1007/s10067-022-06356-z
77. Khanna PP, Gladue HS, Singh MK, et al. Treatment of acute gout: A systematic review. *Semin Arthritis Rheum*. W.B. Saunders. 2014;44(1):31-38. doi:10.1016/j.semarthrit.2014.02.003
78. Schlesinger N. Anti-interleukin-1 therapy in the management of gout. *Curr Rheumatol Rep*. 2014;16. doi:10.1007/s11926-013-0398-z
79. Schlesinger N, Alten RE, Bardin T, et al. Canakinumab for acute gouty arthritis in patients with limited treatment options: Results from two randomised, multicentre, active-controlled, double-blind trials and their initial extensions. *Ann Rheum Dis*. 2012;71:1839-1848. doi:10.1136/annrheumdis-2011-200908
80. Schlesinger N, Pillinger MH, Simon LS, Lipsky PE. Interleukin-1 β inhibitors for the management of acute gout flares: a systematic literature review. *Arthritis Res Ther*. BioMed Central Ltd. 2023;25. doi:10.1186/s13075-023-03098-4
81. Zhang Y, Chen S, Yuan M, Xu Y, Xu H. Gout and Diet: A Comprehensive Review of Mechanisms and Management. *Nutrients*. MDPI. 2022;14. doi:10.3390/nu14173525
82. Dessein PH, Shipton EA, Stanwix AE, Joffe BI, Ramokgadi J. Beneficial effects of weight loss associated with moderate calorie/carbohydrate restriction, and increased proportional intake of protein and unsaturated fat on serum urate and lipoprotein levels in gout: A pilot study. *Ann Rheum Dis*. 2000;59:539-543. doi:10.1136/ard.59.7.539
83. Rai SK, Fung TT, Lu N, Keller SF, Curhan GC, Choi HK. The Dietary Approaches to Stop Hypertension (DASH) diet, Western diet, and risk of gout in men: Prospective cohort study. *BMJ (Online)*. 2017;357. doi:10.1136/bmj.j1794
84. McCormick N, O'Connor MJ, Yokose C, et al. Assessing the Causal Relationships Between Insulin Resistance and Hyperuricemia and Gout Using Bidirectional Mendelian Randomization. *Arthritis and Rheumatology*. 2021;73:2096-2104. doi:10.1002/art.41779
85. Yokose C, McCormick N, Choi HK. The role of diet in hyperuricemia and gout. *Curr Opin Rheumatol*. Lippincott Williams and Wilkins. 2021;33:135-144. doi:10.1097/BOR.0000000000000779
86. Stamostergiou J, Theodoridis X, Ganochoriti V, Bogdanos DP, Sakkas LI. The role of the Mediterranean diet in hyperuricemia and gout. *Mediterr J Rheumatol*. 2018;29:21-25. doi:10.31138/mjr.29.1.21
87. Sacks FM, Campos H. Dietary Therapy in Hypertension. *New England Journal of Medicine*. 2010;362:2102-2112. doi:10.1056/nejmct0911013
88. Juraschek SP, Yokose C, McCormick N, Miller ER, Appel LJ, Choi HK. Effects of Dietary Patterns on Serum Urate: Results From a Randomized Trial of the Effects of Diet on Hypertension. *Arthritis and Rheumatology*. 2021;73:1014-1020. doi:10.1002/art.41614
89. Gee Teng G, Pan A, Yuan JM, Koh WP. Cigarette Smoking and the Risk of Incident Gout in a Prospective Cohort Study. *Arthritis Care Res (Hoboken)*. 2016;68:1135-1142. doi:10.1002/acr.22821
90. Adomako EA, Moe OW. Uric acid transport, transporters, and their pharmacological targeting. *Acta Physiologica*. 2023;238(2):e13980. doi:10.1111/APHA.13980
91. Kang EH, Park EH, Shin A, Song JS, Kim SC. Cardiovascular risk associated with allopurinol vs. benzbromarone in patients with gout. *Eur Heart J*. 2021;42(44):4578-4588. doi:10.1093/eurheartj/ehab619
92. Sanchez-Niño MD, Zheng-Lin B, Valiño-Rivas L, et al. Lesinurad: what the nephrologist should know. *Clin Kidney J*. 2017;10(5):679-687. doi:10.1093/ckj/sfx036
93. Taniguchi T, Ashizawa N, Matsumoto K, et al. Pharmacological Evaluation of Dotinurad, a Selective Urate Reabsorption Inhibitor. *J Pharmacol Exp Ther*. 2019;371(1):162-170. doi:10.1124/jpet.119.259341
94. Sun J, Wang Y, Cheng Y, et al. Efficacy and Safety of Dotinurad Versus Febuxostat for the Treatment of Gout: A Randomized, Multicenter, Double-Blind, Phase 3 Trial in China. *Arthritis Rheumatol*. 2025;77(12):1757-1766. doi:10.1002/art.43261
95. Jun JB, Lee HS, Kim SH, et al. Efficacy and safety of epaminurad, a potent hURAT1 inhibitor, in patients with gout: a randomized, placebo-controlled, dose-finding study. *Arthritis Res Ther*. 2025;27(1):113. doi:10.1186/s13075-025-03577-w
96. Qu Y, Yu Y, Pan J, Li H, Cui C, Liu D. Systematic review and model-based analysis to identify whether

- renal safety risks of URAT1 inhibitors are fully determined by uric acid-lowering efficacies. *Semin Arthritis Rheum.* 2023;63:152279. doi:10.1016/j.semarthrit.2023.152279
97. Shiramoto M, Liu S, Shen Z, et al. Verinurad combined with febuxostat in Japanese adults with gout or asymptomatic hyperuricaemia: a phase 2a, open-label study. *Rheumatology (Oxford).* 2018;57(9):1602-1610. doi:10.1093/rheumatology/kez100
98. Lee HA, Yu KS, Park SI, et al. URC102, a potent and selective inhibitor of hURAT1, reduced serum uric acid in healthy volunteers. *Rheumatology (Oxford).* 2019;58(11):1976-1984. doi:10.1093/rheumatology/kez140
99. Wang Z, Li X, Jin Y, et al. Safety, Efficacy, and Pharmacokinetics of HP501 in Healthy Volunteers and Hyperuricemic Patients: A Phase I/IIa Study. *J Clin Endocrinol Metab.* 2022;107(6):1667-1678. doi:10.1210/clinem/dgac032
100. Cai Z, Lv Y, Liu Y, et al. Quantitative evaluation of clinical efficacy and safety of uric acid-lowering drugs: A modeling analysis of literature aggregate data. *Semin Arthritis Rheum.* 2025;74:152801. doi:10.1016/j.semarthrit.2025.152801
101. Oikonen M, Wendelin-Saarenhovi M, Lyytikäinen LP, et al. Associations between serum uric acid and markers of subclinical atherosclerosis in young adults. The cardiovascular risk in Young Finns study. *Atherosclerosis.* 2012;223(2):497-503. doi:10.1016/j.atherosclerosis.2012.05.036
102. Packer M. Hyperuricemia and Gout Reduction by SGLT2 Inhibitors in Diabetes and Heart Failure: JACC Review Topic of the Week. *J Am Coll Cardiol.* 2024;83(2):371-381. doi:10.1016/j.jacc.2023.10.030
103. Borghi C, Agnoletti D, Cicero AFG, Lurbe E, Virdis A. Uric Acid and Hypertension: a Review of Evidence and Future Perspectives for the Management of Cardiovascular Risk. *Hypertension.* 2022;79(9):1927-1936. doi:10.1161/HYPERTENSIONAHA.122.17956/ASSET/66E5A99E-259D-42A4-84E1-9A71B1A6C2B7/ASSETS/GRAPHIC/HYPERTENSIONAHA.122.17956.FIG04.JPG
104. Watanabe S, Kang DH, Feng L, et al. Uric acid, hominoid evolution, and the pathogenesis of salt-sensitivity. *Hypertension.* 2002;40(3):355-360. doi:10.1161/01.HYP.0000028589.66335.AA/ASSET/107ABD74-700B-4058-9A1A-FFEDC21D18BC/ASSETS/GRAPHIC/24FF5.JPEG
105. Gill D, Cameron AC, Burgess S, et al. Urate, Blood Pressure, and Cardiovascular Disease: Evidence From Mendelian Randomization and Meta-Analysis of Clinical Trials. *Hypertension.* 2021;77(2):383-392. doi:10.1161/HYPERTENSIONAHA.120.16547
106. Wang X, Hou Y, Wang X, et al. Relationship between serum uric acid levels and different types of atrial fibrillation: An updated meta-analysis. *Nutrition, Metabolism and Cardiovascular Diseases.* 2021;31(10):2756-2765. doi:10.1016/J.NUMECD.2021.05.034
107. Lin C, Lai Q, Fang E, et al. Serum urate, cardiovascular mediators, and atrial fibrillation: genetic evidence for URAT1-targeted therapy. *Clin Rheumatol.* 2026;45(2):1401-1411. doi:10.1007/s10067-025-07827-9
108. Pellegrini C, Fornai M, Antonioli L, Blandizzi C, Calderone V. Phytochemicals as Novel Therapeutic Strategies for NLRP3 Inflammasome-Related Neurological, Metabolic, and Inflammatory Diseases. *International Journal of Molecular Sciences* 2019, Vol 20, Page 2876. 2019;20(12):2876. doi:10.3390/IJMS20122876
109. Temmoku J, Fujita Y, Matsuoka N, et al. Uric acid-mediated inflammasome activation in IL-6 primed innate immune cells is regulated by baricitinib. *Mod Rheumatol.* 2021;31(1):270-275. doi:10.1080/14397595.2020.1740410
110. Zhang S, Li D, Fan M, et al. Mechanism of Reactive Oxygen Species-Guided Immune Responses in Gouty Arthritis and Potential Therapeutic Targets. *Biomolecules* 2024, Vol 14, Page 978. 2024;14(8):978. doi:10.3390/BIOM14080978
111. Kang EH, Shin A, Park CS, et al. Risk of urolithiasis associated with allopurinol versus benzbromarone among patients with gout: a population-based cohort study. *Rheumatology (Oxford).* 2024;63(9):2433-2441. doi:10.1093/rheumatology/keae262
112. Qu Y, Yu Y, Pan J, Li H, Cui C, Liu D. Systematic review and model-based analysis to identify whether renal safety risks of URAT1 inhibitors are fully determined by uric acid-lowering efficacies. *Semin Arthritis Rheum.* 2023;63:152279. doi:10.1016/j.semarthrit.2023.152279

This article is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited and it is not used for commercial purposes; 2026, Zehra A. et al., Applied Systems and Discoveries Journals.